

## **RESPONSE OF TWO SOYBEAN CULTIVARS TO DIFFERENT LEVELS ORGANIC FERTILIZER (COMPOST)**

**Abd El-Hafez, G.A.<sup>1</sup> and A. A. Abo El-Soud<sup>2</sup>**

**1-Field Crops Institute, Agric. Res. Center (ARC), Giza, Egypt.**

**2-Microbiology Dept., Soils, Water & Environ. Res. Institute (SWERI), ARC. Giza; Egypt.**

### **ABSTRACT**

Two field experiments were conducted to study the response of two soybean cultivars to different levels of organic fertilizer (compost) on nodulation, nitrogen fixation, growth, yield and yield characters during two successive summer seasons of 2005 and 2006 at the experimental farm of Mallawi Agriculture Research Station, Minia Governorate, Egypt. The soybean cultivars used were Giza 111 and Crawford. Treatments were five different levels of compost, zero, 500, 1000, 1500 and 2000 kg compost/fed compared with the *Bradyrhizobium* inoculated soybean plus starter dose of nitrogen fertilizer (20 kg N/fed) or uninoculated soybean fertilized with the recommended dose of nitrogen fertilizer (70 kg N/fed). A split plot layout with four replications was used.

Results showed that the inoculation with *Bradyrhizobium* in combination with compost gave significant increases in nodule number and dry weight as well as dry weight of shoots and nitrogen content compared with the inoculated plants without compost in Giza 111 and Crawford cultivars.

Also, results showed the superiority of Giza 111 cultivar for increasing nodulation, growth, yield and its components (nodule number and dry weight, shoot dry weight, seed and straw yields/fed, plant height, branches number and pods number per plant, seed weight/plant and 100- seed weight,).

As well as results revealed that all different levels of compost increased significantly growth, seed yield and its components as compared with the control. The highest increase was recorded with 2000 kg rate of compost /fed. followed by 1500 kg/fed. The highest values of plant height, branch number, pods number, seed yield/plant, 100-seed weight, seed and straw yields were obtained from the interaction between Giza 111x 2000 kg compost/fed.

It can be stated that applying different compost levels to soybean plants was the recommended for raising soybean productivity and reducing the environmental pollution under condition of the present study.

### **INTRODUCTION**

Soybean [*Glycine max* (L.) Merrill] is one of the most important annual pulse crops in the world. The cultivated form is used in human food and livestock feeds (Harry and Kwon, 1987). Soybean seeds are one of the main sources of protein and oil in plants. The seed contains approximately 40% protein and 21% edible oil, which used in making margarine. It has been called "yellow jewel", "nature's miracle protein" and "meat of the field" (Noureldin, 1998). The oil is used essentially in margarine, salad oils, cooking oil and shortening. Moreover soybean products became more important because of low costs, nutritionally balanced and beverages for human consumption.

Thus it is important to increase production by increasing the cultivated area or rising per unit area yield by applying the most suitable cultural practices such as compost and growing various new cultivates.

In Egypt, the use of organic materials as fertilizers began to decline, while the use of mineral fertilizers is increasing. At the moment, crop residues are utilized largely for burning, industry and animal feed. The amounts of farmyard

manure available to the Egyptian farmer are not sufficient and it is very poor in organic matter and plant nutrient contents. Hence, the disintegration of the organic fertilizers in soil is very important in order to achieve the important roles of its activity. The available amount of farmyard manure to the Egyptian farmer is not only insufficient but also decreasing with the increasing tendency towards the mechanization of agriculture. An alternative way to meet the growing needs for organic manure is by composting plant and animal residues (Abd-El Ghaffar, 1978). Compost is an eco-friendly fertilizer, it positively improves soil structure, aggregate formation, drought protection, stopping erosion, buffering, reduces fertilizer requirements and gives nutrients when plants need them as well as inoculates the soil with vast numbers of beneficial microbes. Thus, compost can modify soil physical properties and strongly affects its chemical and biological ones (Abdel-Malek *et al.*, 1961; Martin and Gershuny, 1992; Mekail, 1998 and Fontaine *et al.*, 2003).

Many investigators reported that using compost with several crops including legume crops almost duplicated the observed yields besides controlling numerous of soil born diseases (Hoitink *et al.*, 1993). Eghball (2002) reported that after four years of nitrogen or phosphorus based manure and compost applications, soil surface carbon and nitrogen concentration and quantities were greater in the nitrogen compared with the phosphorus based management systems.

Soybean cultivars exhibited differences in its plant height, number of branches, seeds and pods/plant, seed index, seed and straw yields/fed. (Negra and Chirita, 1994 and Abd El-Hafez, 1999).

The aim of this work was to study the effect of different levels of organic compost on the nodulation, growth, yield and yield components for two soybean cultivars plants. Moreover, to minimize the environmental pollution, resulted from the intensive use of chemical fertilization by substituting a part of it with organic fertilizer (compost).

## **MATERIALS AND METHODS**

Two field experiments were conducted during two successive summer seasons of 2005 and 2006 at Mallawi Agric. Res. Station Farm, Minia Governorate, Egypt to study the effect of different levels of organic fertilizer (compost) on nodulation, growth, yield and its components for two soybean cultivars.

### **Soil Sampling:**

Representative soil samples were collected from the top 30 cm layer of the Experimental fields, sieved through 2 mm screen and air-dried. The main physical and chemical properties of the soil in both tested seasons were analyzed according to Page *et al.* (1982) and recorded in Table (1).

### **Microbial strains:**

All microorganisms used in preparing bio-organic fertilizer (compost) such as cellulose decomposers, (*Trichoderma reesei*), phosphate dissolving bacteria (*Bacillus megatherium* var *phosphaticum*) and nitrogen fixing bacteria (*Azotobacter chroococcum*) were prepared in Biofertilizers Production Unit; Microbiology Department; SWERI, ARC, Giza, Egypt. Also, mixtures of two strains of *Bradyrhizobium japonicum* USDA 110 and HH303 (approximately 10<sup>9</sup> cells/ml) used in soybean inoculation were prepared and added to sterile solid carrier (vermiculite + 10% peat) to prepare the inoculant used for soybean inoculation.

**Table (1): Some physical and chemical properties of the surface layer (0.0-30cm) of the studied soil at two seasons of 2005 and 2006**

Physical and chemical properties	First season 2005	Second season 2006
<b>Particles size distribution</b>		
Sand %	20.70	19.95
Silt %	34.05	35.00
Clay %	45.25	45.05
Texture grade	Clayey	Clayey
pH (1:5 soil water suspension)	8.00	7.92
E.C. (dS/m at 25 °c)	1.75	1.61
Organic matter %	0.94	0.96
<b>Soluble cations (meg/L)</b>		
Ca <sup>++</sup>	4.27	4.14
Mg <sup>++</sup>	2.34	2.41
Na <sup>+</sup>	7.49	7.46
K <sup>+</sup>	2.72	2.65
<b>Soluble anions (meg/L)</b>		
CO <sub>3</sub> <sup>-</sup>	0.00	0.00
HCO <sub>3</sub> <sup>-</sup>	1.86	1.91
Cl <sup>-</sup>	2.39	2.41
SO <sub>4</sub> <sup>-</sup>	12.57	12.34

**Preparation of compost:**

Compost was a mixture of farm residues (corn and cotton stalks, sesame and soybean straw) inoculated with special bio-decomposer strains of bacteria, fungi under aerobic condition. First, enriched compost was prepared using raw materials of farm residues shredded into 1.5 cm. Farmyard manure collected from Mallawi, Agric. Res. Station farm, Minia Governorate, Egypt were used. Moreover, some microbial inoculants such as cellulose decomposers, phosphate dissolving bacteria and nitrogen fixing bacteria were added to the mixture.

To set up the heap, the remain raw materials were mixed with farmyard manure and inoculated with microbial then built in successive layers tamped well over the bed mixture of raw materials up to 1m high. Moisture was maintained to 60% and the moistening was considered satisfactory when a hand full of composted materials would wet the hand but not drip. This heap was turned up down every 15 days for 75 days (till maturation). Samples were taken at maturation of heap construction, mixed thoroughly, air dried and ground to chemical, physical and microbiological determinations (Table 2).

**Plant cultivars:**

Soybean seeds Giza 111 and Crawford were kindly provided by Field Crops Research Institute, ARC, Giza, Egypt.

**Field experiments:**

Two consequence field experiments were done at the experimental farm of Mallawi Agriculture Research station. Minia, Governorate, Egypt.

The compost was added before cultivation during soil preparation. A split plot design was used in this study with four replicates. The tested cultivars were set up randomly in the main plots while the compost treatments were arranged at random in subplots of 3x3.5 m with 60 cm apart between ridges (5 ridges/subplot). Compost was added at the rates of zero, 500, 1000, 1500 and 2000 kg/fed at 15 days before cultivation during soil preparation. Super-phosphate (15.5% P<sub>2</sub>O<sub>5</sub>) and potassium sulfate (48% K<sub>2</sub>O) were incorporated into soil before sowing at the rates of 200 and 50 kg/fed, respectively.

**Table (2): Characteristics the bio-compost used in the field experiment**

Properties	Values
pH	7.28
E.C. (dS/m at 25°C)	4.18
Organic matter (%)	36.07
Organic-C (%)	24.97
Total N ( % )	1.20
C/N ratio	20.8
Total-P (%)	1.04
Total-K (%)	0.52
Total soluble-N (ppm)	642.5
Available-P ( ppm)	276.0
Available-K (ppm)	725.4
Total count of bacteria	26 x 10 <sup>7</sup>
Total count of fungi	16 x 10 <sup>6</sup>
Total count of actinomycetes	10 x 10 <sup>6</sup>

Sowing soybean seeds took place on May 4<sup>th</sup> and 10<sup>th</sup> during summer seasons of 2005 and 2006, respectively. Soybean seeds were cultivated as recommended practice and inoculated with effective strains of *Bradyrhizobium japonicum* using Arabic gum solution as adhesive material just before sowing. Inoculant was applied at rate of 600 g per 60 kg soybean seeds. Soybean seedlings were thinned out to two plants/hill at 21 days from sowing. The other recommended agronomic practices for soybean cultivation were used just before sowing. The control treatments were inoculated with soybean plus starter dose of nitrogen fertilizer (20 kg N/fed) and uninoculated fertilized with the recommended dose of nitrogen fertilizer (70 kg N/fed).

**The following treatments were studied:**

- Bradyrhizobium* inoc. + 0 Kg compost/fed
- Bradyrhizobium* inoc. + 500 Kg compost/fed
- Bradyrhizobium* inoc. + 1000 Kg compost/fed
- Bradyrhizobium* inoc. + 1500 Kg compost/fed
- Bradyrhizobium* inoc. + 2000 Kg compost/fed
- Bradyrhizobium* inoc. + 20 Kg N/fed
- Un-inoc. + 70KgN/fed

Five plants samples were randomly taken from each plot at 60 days after planting to determine nodule number and dry weight as well as shoot dry weight and its nitrogen contents according to Page *et al.* (1982)...

At harvest, ten guarded plants were randomly chosen to determine plant height (cm) number of branches and pods/ plant, seed weight (g) /plant and weight of 100-seed (seed index). Also, seed and straw yield (ton/fed) were estimated from the three central ridges of each subplot. Seed and straw nitrogen content were determined according to Page *et al.* (1982).

**Statistical analysis:**

The analysis of variances was made separately for each season, and then a combined analysis of seasons, cultivars and treatments was made according to Gomez and Gomez (1984).

**RESULTS AND DISCUSSION**

**1-Effect of the interaction:**

Results of two soybean cultivars after 60 days of plant grown under two successive seasons as affected by *Bradyrhizobium* inoculation combined with

different levels of organic fertilizer (compost) are given in Table (3). Obtained data showed that all nodulation and vegetative growth parameters were significantly affected by different treatments under study. The uninoculated plants received the recommended dose of 70 kg N/fed did not produce nodules number or dry weight in the two cultivars at both seasons, indicating that the specific indigenous *Bradyrhizobium japonicum* in the experimental soils are absent.

**Table (3): Effect of *Bradyrhizobium* inoculation combined with different levels of compost on nodulation and growth of two soybean cultivars during the two seasons of 2005 and 2006**

Treatments	Nodulation				Shoot			
	Number / plant		Dry weight (mg/plant)		Dry weight (g/plant)		N-content (mg/plant)	
	Giza 111	Crawford	Giza 111	Crawford	Giza 111	Crawford	Giza 111	Crawford
<b>Season 2005</b>								
Inoc. + 0 Kg compost/fed	8	6	86.2	72.3	28.5	21.6	584.3	399.6
Inoc.+500 Kg compost/fed	15	10	163.5	98.4	35.4	32.6	764.6	625.9
Inoc.+1000 Kg compost/fed	19	12	243.6	102.6	38.6	35.7	907.1	731.9
Inoc. + 1500 Kg compost/fed	20	15	257.4	115.4	48.5	42.8	1290.1	1005.8
Inoc. + 2000 Kg compost/fed	22	16	268.4	187.6	56.2	46.8	1539.9	1132.6
Inoc. + 20 Kg N/fed	20	18	254.8	151.6	51.2	44.6	1318.6	1052.6
Un-inoc. 70KgN/fed	0	0	0	0	54.7	49.2	1559.0	1254.6
LSD 0.05	2.1		20.6		5.3		186.4	
<b>Season 2006</b>								
Inoc. + 0 Kg compost/fed	10	8	92.5	84.6	30.4	27.6	656.64	524.4
Inoc. + 500 Kg compost/fed	18	13	186.3	105.6	36.5	33.5	448.95	643.2
Inoc. + 1000Kg compost/fed	22	16	266.5	113.5	46.8	36.8	1132.56	721.28
Inoc. + 1500Kg compost/fed	25	18	2648	135.9	50.2	44.3	1345.36	877.14
Inoc. + 2000Kg compost/fed	27	20	292.6	165.7	53.4	48.5	1452.48	1028.2
Inoc. + 20 Kg N/fed	26	18	284.7	157.6	52.6	45.8	1378.12	893.1
Un-inoc.+70Kg N/fed	0	0	0	0	55.8	49.6	1551.24	1066.4
LSD 0.05	2.6		34.6		6.2		198.6	

Inoculation with *Bradyrhizobium japonicum* plus starter dose of nitrogen caused significant increases over the inoculated without nitrogen fertilizer control in number & dry weight of nodules by 150.0 & 195.6% for Giza 111 cultivar and by 200.0 & 109.7% for Crawford cultivar in the first season, respectively. The corresponding increases in the second season were 160.0 & 207.8% and 200.0 & 86.3%, respectively.

Moreover, *Bradyrhizobium* inoculation combined with compost (2000 kg compost/fed) caused significant increases in nodule number and dry weight compared to *Bradyrhizobium* inoculation alone. These increases at the first and second seasons were 175.0 and 170.0% in Giza 111 cultivar, respectively. The corresponding increases in Crawford cultivar were 166.7 and 150.0% in the same order during two seasons. The increases in nodules dry weight due to these treatments over the inoculation without compost in Giza 111 variety were 211.4, 216.3% during the first and second seasons, respectively. While, these increases were 159.5 and 95.9% for Crawford variety at first and second seasons, respectively.

Data presented in Table (3) revealed that inoculated treatment without compost recorded the lowest dry weight of Giza 111 shoots (28.5 and 30.4 g/plant) and the lowest nitrogen content (584.4 and 656.6 mg N/plant) at the first and second season, respectively.

Similar trends were obtained in inoculated Crawford cultivar without compost, where shoots dry weight recorded 21.6 & 27.6 g/plant and nitrogen content of 399.6 & 524.4 mg N/plant during the two seasons in the same order.

In Giza 111 cultivar, the uninoculated plants received the full dose of nitrogen (70 kg N/fed) increased shoot dry weight & nitrogen content by 91.9 & 166.8% and 83.6 & 136.2% over the control at the first and second season, respectively. While, in Crawford cultivar these increases were 127.8 & 214.0% and 79.7 & 103.4% in the same order during the two seasons.

Inoculation with *Bradyrhizobium* in combination with compost gave significant increases in dry weight of shoots and nitrogen content compared with the inoculated plants without compost in Giza 111 and Crawford cultivars.

*Bradyrhizobium* in combination with compost (2000 kg compost/fed) increased shoots dry weight & N-content of Giza 111 plant by 97.2 & 163.5% over the *Bradyrhizobium* alone, respectively at the first season. These increases were 75.7 and 121.2% at the second season. While, in Crawford cultivar the increases in dry weight of shoots & N-content were 116.4 & 183.4% in the same order at the first season. In the second season, these increases were 75.7 & 96.1%, respectively.

These increases could be due to the biological role of *Rhizobium* in enhancing plant growth and N<sub>2</sub>-fixation as reported by Abo El-Soud *et al.* (2004) and Mekhemar *et al.* (2005).

Data again revealed that the maximum dry weight of shoots and nitrogen content were obtained in plants inoculated with *Bradyrhizobium* combined with 2000 kg compost/fed followed by those inoculated with *Bradyrhizobium* combined with starter dose of nitrogen fertilizer.

The interaction between two soybean cultivars and different levels of compost on the plant high, branch number, pods number per plant and 100-seed weight in 2005 and 2006 seasons are presented in Table (4). The highest values (118.0 cm, 3.3 branches, 7.9 pods and 20.9 g, respectively) were obtained by sowing Giza111 cultivar with 2000 kg compost/fed followed by, sowing Giza111 with 1500kg compost /fed, while sowing Crawford with zero (control) recorded the lowest in all previous mentioned characters.

Regarding seed and straw yields of Giza111 and Crawford cultivars, results in Table (5) exhibited significant differences in seed and straw yields between plants inoculated with *Bradyrhizobium* alone or combined with different compost levels and uninoculated plants, which received 70 kg N/fed. The inoculation of plants without compost application recorded the lowest values of seed and straw yields in the first and second seasons. The highest seed yield was achieved with 2000 kg compost/fed during both tested seasons. Then yield tended to decrease due to the use of 1500 kg compost/fed. The significant differences in seed and straw yields were recorded as compost levels from zero to 500,1000.1500 and 2000 kg/fed and soybean cultivars. Soybean, Giza 111 inoculated with *Bradyrhizobium* combined with 500, 1000, 1500 and 2000 kg compost/fed increased seed yield by 32.1, 74.5, 130.5 and 196.3% and straw yield by 38.0, 72.4, 159.8 and 166.6% over the inoculated plants without compost in the first season, respectively. While, in the second season which recorded



increases of 30.1, 66.6, 117.0 and 123.6 % in seed yield at the same order for plants inoculated with *Bradyrhizobium* combined with 500, 1000, 1500 and 2000 kg compost/fed. The increases in straw yield were 11.6, 40.0, 75.9 and 82.0%, respectively.

Similar trends were observed in Crawford cultivar, when inoculated with *Bradyrhizobium* combined with 500, 1000, 1500 and 2000 kg compost/fed, since the seed yield increased by 25.6, 61.0, 84.4 and 109.5% and straw yield by 11.7, 45.8, 60.5 and 96.7% over the inoculated plants without compost in the first season, respectively. While, in the second season, which recorded increases of 14.8, 23.3, 59.5 and 63.2% in seed yield at the same order for plants inoculated with *Bradyrhizobium* combine with 500, 1000, 1500 and 2000 kg compost/fed. The increases in straw yield were 11.6, 40.0, 75.9 and 82.0%, respectively.

The increase in seed and straw yields of soybean cultivars may be due to that compost has direct impacts on plant growth, yield and its attributes; these direct impacts come from providing the plants by the important nutrients for growth and metabolism. Adjacent to the direct impacts, there are two indirect ones, the first is out of compost hydrolysis in soil, and humic acid is produced, which played an important role in reducing soil pH and increasing soil nutrient availability to the plants. Beside the role of humic acid as a rich fertilizer itself, the second indirect impact is the preferable consequence of compost on the soil mechanical properties to improve soil physical properties including water-holding capacity and reduces the probability of soil borne infection (Hoitink *et al.*, 1993).

**Table (4): Effect of *Bradyrhizobium* inoculation combined with different levels of compost on yield component of two soybean cultivars during the two seasons of 2005 and 2006**

Treatments	Plant height (cm)		Branches number /plant		Pods number / plant		100-Seed weight (g)	
	Giza 111	Crawford	Giza 111	Crawford	Giza 111	Crawford	Giza 111	Crawford
<b>Season 2005</b>								
Inoc. + 0 Kg compost/fed	62.7	58.3	1.3	1.2	36.0	30.3	11.5	10.4
Inoc.+500 Kg compost/fed	75.8	63.7	1.7	1.5	45.3	35.3	13.4	11.3
Inoc.+1000 Kg compost/fed	88.0	74.6	2.6	2.5	53.1	40.7	14.9	12.3
Inoc. + 1500 Kg compost/fed	109.4	81.1	3.2	2.8	63.2	49.3	19.7	15.1
Inoc. + 2000 Kg compost/fed	118.0	84.2	3.3	2.8	71.9	44.7	20.9	15.3
Inoc. + 20 Kg N/fed	112.6	79.2	2.8	2.8	58.8	45.2	18.7	14.8
Un-inoc.+ 70KgN/fed	114.5	89.6	3.4	3.1	69.8	46.8	20.8	16.7
LSD 0.05	7.1		0.3		5.0		2.1	
<b>Season 2006</b>								
Inoc. + 0 Kg compost/fed	62.4	53.3	1.3	1.0	41.3	33.3	12.8	11.5
Inoc. + 500 Kg compost/fed	73.8	60.3	1.9	1.3	47.7	36.7	14.4	12.7
Inoc. + 1000Kg compost/fed	83.0	70.3	2.9	1.9	56.4	42.7	16.1	13.6
Inoc. + 1500Kg compost/fed	100.7	75.2	3.4	2.2	70.9	52.3	20.2	16.1
Inoc. + 2000Kg compost/fed	108.0	76.3	3.8	2.8	78.3	46.3	22.6	15.7
Inoc. + 20 Kg N/fed	98.5	74.6	3.2	2.4	68.9	45.2	18.6	15.2
Un-inoc.+70Kg N/fed	104.0	78.4	3.5	3.0	74.5	50.6	22.5	16.8
LSD 0.05	6.4		0.7		6.1		4.0	

From the abovementioned results, it can be stated that the application of 2000 kg compost/fed before sowing soybean could be recommended for raising soybean productivity and decreasing pollution through decreasing mineral nitrogen application under the environmental condition of the present study.

**Table (5): Effect of *Bradyrhizobium* inoculation combined with different levels of compost on seed and straw yield of two soybean cultivars during the two seasons of 2005 and 2006**

Treatments	Yield (Ton/fed)				%N			
	Seed		Straw		Seed		Straw	
	Giza 111	Crawford	Giza 111	Crawford	Giza 111	Crawford	Giza 111	Crawford
<b>Season 2005</b>								
Inoc. + 0 Kg compost/fed	1.070	0.860	1.143	1.095	3.12	2.82	1.54	1.30
Inoc.+500 Kg compost/fed	1.413	1.080	1.577	1.223	3.20	3.14	1.65	1.39
Inoc. + 1000 Kg compost/fed	1.867	1.384	1.970	1.597	3.20	3.12	1.86	1.50
Inoc. + 1500 Kg compost/fed	2.457	1.586	2.970	1.757	3.32	3.15	2.36	1.84
Inoc.+2000 Kg compost/fed	3.033	1.802	3.047	2.143	3.64	3.24	2.41	2.20
Inoc. + 20 Kg N/fed	2.275	1.420	2.859	1.546	3.42	3.22	2.40	2.00
Un-inoc. + 70KgN/fed	2.937	1.872	3.165	2.420	3.65	3.48	2.50	2.29
LSD 0.05	0.320		0.436		0.70		0.51	
<b>Season 2006</b>								
Inoc. + 0 Kg compost/fed	1.163	1.011	1.323	1.150	3.16	2.88	1.62	1.42
Inoc. + 500 Kg compost/fed	1.513	1.161	1.893	1.283	3.28	3.14	1.76	1.49
Inoc. + 1000Kg compost/fed	1.937	1.247	2.063	1.610	3.30	3.12	1.94	1.65
Inoc. + 1500Kg compost/fed	2.524	1.613	3.023	2.023	3.43	3.15	2.44	1.85
Inoc. + 2000Kg compost/fed	2.600	1.650	3.187	2.093	3.68	3.24	2.49	1.95
Inoc. + 20 Kg N/fed	2.485	1.425	2.435	1.985	3.52	3.22	2.20	2.26
Un-inoc.+70Kg N/fed	2.678	1.843	3.285	2.135	3.70	3.48	2.29	2.32
LSD 0.05	0.401		0.485		0.75		0.61	

**2- Statistical main effect of compost:**

Regardless of seasons and cultivars, results in Table (6) indicated that the highest means of nodules number and dry weight, shoot dry weight and its nitrogen content as well as seed and straw yields were obtained in plants inoculated with *Bradyrhizobium* and fertilized with 2000 kg compost/fed. There are insignificant difference between the use of 2000 kg compost/fed and/or the use of 20 kgN/fed.

Table (6) shows that the maximum number of nodules and their dry weight were obtained in plants inoculated with *Bradyrhizobium* combined with 2000 kg compost/fed followed by those inoculated with *Rhizobium* plus 20 kg nitrogen/fed and those inoculated with *Bradyrhizobium* plus 1500 kg compost/fed. This could be due to the effect of the applied compost levels on plant growth, which is resulting from the production of plant growth regulators, vitamins leading to enhancement the uptake of plant nutrients, suppression of pathogenic of deleterious organisms as reported by many worker (Chebotar *et al.*, 2001; Zaid *et al.*, 2003 and Kennedy *et al.*, 2004). Irrespective of inoculation, the nodulation exhibited a significant increase due to the application of compost at different levels in comparison to the treatments received no compost. Data also revealed significant increases of nodules number and dry weight for seed inoculated with



*Bradyrhizobium* compared to uninoculated ones. In fact, the addition of enriched organic fertilizer to soil led to raise its fertility and microbial activity, which reflected on enhancing the root proliferation and nodule formation. Such promotion nodulation pattern for many legumes is confirmed by many investigators (El-Sawi *et al.*, 2001; Bai *et al.*, 2002; Abdel-Wahab and Ahmed, 2003 and Abdel-Wahab and Said, 2004).

Concerning the dry weight of 60 day-old soybean plants, data in Table (6) clearly illustrated that the addition of organic fertilizer led to a higher increase in shoots dry weight and its nitrogen content to be similar or higher than values obtained by using the recommended treatment. As well as, plant height, branches and pods number/ plant, seed index (Table 7) seed and straw yield, its nitrogen percentage (Table 8). This positive effect was magnified when combined with *Bradyrhizobium* inoculation particularly at high level of compost (1500 and/ or 2000 kg/fed), which gave the highest values of these parameters. Using such treatments exhibited the increases in characters under study over the control (zero compost). There are no significant differences due to the use any of 2000 kgN/fed, 1500 kg/fed or fertilizer with starter dose of 20 kg N/fed, respectively.

**Table (6): Statistical main effect of *Bradyrhizobium* inoculation combined with different levels of compost on nodulation and growth of soybean during the two seasons of 2005 and 2006**

Treatments	Nodulation		Shoot	
	Number / plant	Dry weight (mg/plant)	Dry weight (g/plant)	N-content (mg/plant)
Inoc. + 0 Kg compost/fed	8.0	83.9	27.0	541.2
Inoc. + 500 Kg compost/fed	14.0	138.5	34.5	620.7
Inoc. + 1000 Kg compost/fed	17.3	181.6	39.5	873.2
Inoc. + 1500 Kg compost/fed	19.5	193.4	46.5	1129.6
Inoc. + 2000 Kg compost/fed	21.3	228.6	51.2	1288.3
Inoc. + 20 Kg N/fed	20.5	212.2	48.6	1160.56
Un-inoc. + 70KgN/fed	0.0	0.0	52.3	1357.8
LSD 0.05	2.1	20.4	4.5	84.2

**Table (7): Statistical main effect of *Bradyrhizobium* inoculation combined with different levels of compost on yield component of soybean during the two seasons of 2005 and 2006**

Treatments	Plant height	Branches number /plant	Pods number /plant	100-Seed weight (g)
Inoc. + 0 Kg compost/fed	59.18	1.2	35.225	11.55
Inoc. + 500 Kg compost/fed	68.40	1.6	41.25	12.95
Inoc. + 1000 Kg compost/fed	78.98	2.475	48.225	14.225
Inoc. + 1500 Kg compost/fed	91.6	2.9	58.925	17.775
Inoc. + 2000 Kg compost/fed	96.63	3.175	60.3	18.625
Inoc. + 20 Kg N/fed	91.23	2.8	54.525	16.825
Un-inoc. + 70KgN/fed	966.25	3.25	60.425	19.2
LSD 0.05	5.3	0.4	4.8	3.2

The increases in the previous parameters may be attributed in part to the effect of organic fertilizer used on the production of humus substances, which improves the physical and chemical properties of soil, increases the water holding capacity, availability of nutrients, which leads to establish suitable growth media for growing plants. Claims to the promotion effect of enriched bio-organic soil conditioner on plant growth were reported by various workers, Bashan and Levanony (1990), Antoun *et al.* (1998) and Abdel-Wahab and Ahmed (2003) who explained the favorable effects of the combination between compost and biofertilizers on the basis of the beneficial effects of bacteria on the nutrient

availability, vital enzymes, hormonal stimulating effects on plant growth or the increasing of photosynthetic activity.

**Table(8):Statistical main effect of *Bradyrhizobium* inoculation combined with different levels of compost on seed and straw yields of soybean during the two seasons of 2005 and 2006**

Treatments	Yield (ton/fed)		%N	
	Seed	Straw	Seed	Straw
Inoc. + 0 Kg compost/fed	1.026	1.178	2.995	1.470
Inoc. + 500 Kg compost/fed	1.292	1.494	3.190	1.573
Inoc. + 1000 Kg compost/fed	1.609	1.810	3.185	1.738
Inoc. + 1500 Kg compost/fed	2.045	2.443	3.263	2.123
Inoc. + 2000 Kg compost/fed	2.271	2.620	3.450	2.263
Inoc. + 20 Kg N/fed	1.901	2.206	3.345	2.215
Un-inoc. + 70KgN/fed	2.333	2.751	3.578	2.350
LSD 0.05	0.280	2.310	0.520	0.492

### 3- Statistical main effect of cultivars

Data in Table (9) showed that the application of different levels of compost combined with *Bradyrhizobium* inoculation showed significant differences between the soybean cultivars due to all studied characters in both seasons except for seed and straw N %, nodules number. Giza111 cultivar exceeded Crawford cultivar in nodules number and plant dry weight, growth, plant height, branches and pods number, seed index, seed and straw yields.

Such differences between cultivars might be attributed to the growth habit of each cultivar, which is governed by genetically factors and/or environmental condition. These results are in agreement with those reported by Zhou *et al.* (1994), El-Karamity (1998) and Abd El-Hafez (1999).

### 4- Statistical main effect of season:

The analysis of variance for seasons, treatments and their interactions of all studied characters are given in Table (9). Data revealed that there are significant differences in plant height, dry weight of shoots and shoots N-content as well as seed and straw yields.

**Table (9): Response of soybean parameters due to Statistical main effect of two cultivars and two seasons over *Bradyrhizobium* inoculation combined with different levels of compost**

Growth parameters	Cultivars			Seasons		
	Giza 111	Crawford	LSD 0.05	2005	2006	LSD 0.05
Nodules number per plant	13.6	10.0	1.6	12.9	15.8	1.4
Nodules DW (mg/plant)	156.5	87.7	17.5	143.0	153.6	14.8
Shoot DW (g/plant)	37.5	32.9	2.4	41.7	43.7	1.8
Shoot N-content (mg/plant)	937.0	703.3	145.2	1011.9	979.9	126.8
Plant height (cm)	78.0	60.0	6.2	86.9	80.6	4.4
Number of branches per plant	2.3	1.8	0.2	2.5	2.5	0.1
Number pods per plant	49.2	35.3	4.2	49.3	53.2	3.4
100-Seed weight (g)	14.5	11.6	1.8	15.4	16.3	1.3
Seed yield ton/fed	1.76	1.17	0.22	1.790	1.775	0.12
Straw yield ton/fed	2.00	1.42	0.45	2.037	2.106	0.12
%N in seed	2.80	2.26	0.36	3.265	3.307	0.23
%N in straw	1.73	1.50	0.22	1.945	1.977	0.18

The second season of 2006 recorded the highest values of nodule number (15.8), nodule dry weight (153.6), shoot dry weight (43.7g/plant) as well

as, the highest values of straw yield 2.106 ton/fed in the second season. While, in the first season of 2005 the highest values of shoots N-content (1011.9 mg N/plant), plant height (86.9cm) and seed yield (1.790 ton/fed) were recorded. These results indicated that season effect is important in the study. The differences between seasons reflecting the differences in weather and the environmental conditions along both studied seasons, suggesting the possibility to rise yield level by choosing the proper time of agriculture and other agronomic practices.

## REFERENCES

- Abd El-Ghaffar, A.S. (1978). The significance of organic materials to Egyptian agriculture and maintenance of soil productivity. FAO/STDA workshop on organic materials and soil productivity in the Near East 9-18 Oct.,1978, Alex., Egypt.
- Abd El-Hafez, G. A. (1999). Studied the effect of sowing date and plant population density on growth and yield of some soybean cultivars, Ph.d. Thesis, Fac. Agric., Minia Univ., Egypt.
- Abdel-Malek, Y. ; M. Monib and M.N. Zayed (1961). Bacteriological and chemical studies in rice straw compost. J. Soil Sci., U.R.E., 1: 51-66.
- Abdel-Wahab, A.F.M. and A.S. Ahmed (2003). Preparation of bio-organic materials or their utilization as soil amendment and growth media. Annals Agric. Sci., Ain Shams Univ., Cairo, 48: 561-572.
- Abdel-Wahab, A.F.M. and M.S. Said (2004). Response of Faba bean to bio and organic fertilization under calcareous Soil Conditions Egypt. J. Appl. Sci; 19 (1) 305-320.
- Abo El-Soud, A.A.; M.A. El-Deeb and Kh.M. El-Yamani (2004). Improving of faba bean production by application of organic compost and rhizobial inoculation under newly reclaimed soil. Egypt. J. Appl. Sci., 19: 333-344.
- Antoun, H.; J. Chantal.; N. Goussard.; R. Chabot and R. lalande (1998). Potential of *Rhizobium* and *Bradyrhizobium* species as plant growth promoting rhizobacteria on non-legumes: Effect on radishes (*Raphanus sativus* L.) Plant and Soil. 204 : 57-67.
- Bai, Y., Pan, B., Charles, T.C. and Smith, D. L. (2002). Co-inoculation dose and root zone temperature for plant promoting rhizobacteria on soybean [*Glycine max* (L.) Merr] grown in soil less media. Soil Biology and Biochemistry, 34:1953-1957.
- Bashan, Y and H. Levanony (1990). Current status of *Azospirillum* inoculation technology : *Azospirillum* as a challenge for agriculture. Can. J. Microbiol. 36: 591-608.
- Chebotar, V.K.; C.V. Asis Jr and S. Akao (2001). Production of growth-promoting substances and high colonization ability of rhizobacteria enhance the nitrogen fixation of soybean when co-inoculated with *Bradyrhizobium japonicum*. Biol. Fertil. Soils, 34: 427-432.
- Eghball, B. (2002). Soil properties as influenced by phosphorus and nitrogen-based manure and compost applications. Agron. J. 94: 128-135.
- El-Karamity, A.E. (1998). Performance of soybean cultivars at different soil moisture levels. Zagazig. Agric. Res., 25(2):195-210.
- El-Sawi, M. M., A. F. Abdel-Wahab and A. A. Abo El-Soud (2001). Effect of commercial biofertilizer "Nemaless" on nodulation, growth and yield of faba bean. J. Agric. Sci. ansoura univ., 26: 3201 -3o26.
- Fontaine, S ;A. Mariotti and L.Abbadie(2003).he priming effect of organic matter:a question of microbial competition. Soil Biol.&Biochem.,35: 837-843.
- Gomez, K.A. and A.A. Gomez (1984). Statistical Procedures for Agricultural Research. 2<sup>nd</sup> edition. John Willey and Sons, New York.

- Harry, E.S. and T.W. Kown (1987). Soybean utilization. Van. Nostrand Reinhold Company, New York, pp. 346.
- Hoitink, H.; M. Boehm and Y. Hadar (1993). Mechanisms of suppression of soil borne plant pathogens in compost amendment substrats. pp. 601-621. In: H. Hoitink and H. Keener (eds.) Science and Engineering of Composting Renaissance Publications, Worthington. OH.
- Kennedy, I.R.; A.T.M.A. Choudhury; Mihaly L. Kecskes(2004). Non-symbiotic bacterial diazotrophs in crop-farming system: can their potential for plant growth promotion be better exploited Soil Biol. And Biochem, 36:1229-1244.
- Martin, D.L. and G. Gershuny (1992). "The Rodale Book of Composting". Rodale Press, Emmaus, Pennsylvania.
- Mekail, M.M. (1998). Evaluation of some natural organic wastes as amendments for virgin coarse textured soils. I. Effect of filtermud (Pressmud) and nitrogen application on some soil properties and wheat yield. J. Agric. Sci., Mansoura Univ., 23: 5749-5762.
- Mekhemar, G.A.A.; M. Shaaban; A.A. Ragab and A.M.M. Biomy (2005). Response of fab bean to inoculation with *Rhizobium leguminosarum* bv. *Viciae* and plant growth promoting rhizobacteria under newly reclaimed soils. Egypt. J. Appl. Sci., 20: 126-144.
- Negra, S. and N. Chirita (1994). Response of some new soybean cultivars and lines under conditions of the Agric., Res., Station at Secuiem Cecetari Agronomic in Moldova 27 (1-2): 121-126.
- Noureldin, N.A. (1998). The History and Development of Soybean as a Crop in Egypt, Proceedings of the International Conference on Soybean Production under Newly Reclaimed Lands in Egypt, November, 28-29.
- Page, A.I.; R.H. Miller and D.R. Keeney (1982). Methods of Analysis. Part 2, Chemical and Microbiological Properties, Second Edition, Amr. Soc. Agron. Inc., Amr. Soil Sci. Soc. pp. 595-624.
- Zaied, K.A.; A.H. Abd El-Hady; Aida H. Afify and M.A. Nassef (2003). Yield and nitrogen assimilation of winter wheat inoculated with new recombinant inoculants of rhizobacteria. Pakistan J. Biol. Sci., 6: 344-358.
- Zhou, Z.Q.; N. Xu; Z.Y. Liu; L.J. Wang and Y.Q. Wang (1994). A study of agronomic characters and yield formation in summer sown soybean cultivars in Northeastern Hebei Soybean Sci. 12:340-346.

### استجابة صنفان من فول الصويا لمستويات مختلفة من السماد العضوي (الكمبوست).

جمال عبد العزيز عبد الحافظ<sup>1</sup> - علاء الدين عبد الحميد أبو السعود<sup>2</sup>  
1- معهد بحوث المحاصيل الحقلية - مركز البحوث الزراعية - الجيزة - مصر  
2- معهد بحوث الأراضي والمياه والبيئة - مركز البحوث الزراعية - الجيزة - مصر

أجريت تجربتان حقليتان موسمي 2005-2006 بالمزرعة البحثية بمحطة البحوث الزراعية بملوى محافظة المنيا لدراسة استجابة صنفان من فول الصويا جيزة 111، كراوفورد للتلقيح بالريزوبيا في وجود مستويات مختلفة من السماد العضوي (الكمبوست) 0، 500، 1000، 1500، 2000 كجم كمبوست/فدان بالإضافة لمعاملات المقارنة (التلقيح بالريزوبيا مع إضافة جرعة تنشيطية 20 كجم ن/فدان من السماد النيتروجيني أو التسميد المعدني الموصى به 70 كجم ن/فدان) وذلك في أربعة مكررات واستخدم نظام القطع المنشقة مرة واحدة في التصميم التجريبي حيث وزعت معاملات الأصناف في القطع الرئيسية ومعاملات التسميد الحيوي (الكمبوست) في القطع المنشقة الثانوية. وكانت أهم النتائج المتحصل عليها كما يلي .

أدى التلقيح بالريزوبيا في وجود الكمبوست إلى زيادة في أعداد العقد الجذرية وأوزانها وكذلك المجموع الخضري ومحتواه من النيتروجين بالمقارنة بالتلقيح بدون إضافة الكمبوست في الصنفين تحت الدراسة. أظهر صنف جيزة 111 تفوق في كل الصفات المختبرة وهي (أعداد وأوزان العقد الجذرية - الوزن الجاف للنبات - طول النبات عدد الفروع /نبات - عدد القرون على النبات، وزن ال100 بذرة ووزن محصولي البذور والقش. أشارت النتائج إلى أن زيادة مستويات التسميد العضوي (الكمبوست) أدت إلى زيادة جميع الصفات موضع الدراسة مقارنة بالكنترول. وقد حقق مستوى التسميد العضوي 2000 كجم/فدان تفوقاً على جميع المستويات في كل الصفات موضع الدراسة تلاه مستوى التسميد 1500 كجم/فدان.

كان للتفاعل بين العوامل تحت الدراسة تأثيراً إيجابياً حيث سجلت أعلى قيم (طول النبات - عدد الفروع/نبات - عدد القرون /نبات، وزن محصول البذور والقش طن/الفدان) لذلك ينصح بزراعة الصنف جيزة 111 مع التسميد العضوي (الكمبوست) 2000 كجم/فدان ويمكن أن نستخلص من النتائج أن معاملة نباتات فول الصويا بالتسميد العضوي في وجود التلقيح بالريزوبيا يزيد من إنتاجية الفدان وكذلك يقلل من التلوث البيئي الناتج عن الاستخدام المكثف للأسمدة المعدنية.