

# Journal of Agricultural Chemistry and Biotechnology

Journal homepage & Available online at: [www.jacb.journals.ekb.eg](http://www.jacb.journals.ekb.eg)

## Bioremoval of Lead by *Bacillus amyloliquefaciens* Mpa 1034 Isolated from Wastewater Textile Factories at El-Mahala El-Kobra City, Egypt

Mai M. A. Elgogary; M. A. E. Selim and Eman H. Ashour\*

Microbiology Department, Faculty of Agriculture, Mansoura University, Egypt



Cross Mark



### ABSTRACT

Heavy metals are considered serious environmental pollutants. Industrialization activity has put high amounts of toxic effluents, containing toxic metals into the environment. Therefore, there is an urgent need for industrial wastewater treatment. This study aims to lead bioremoval by active Pb-resistance bacterial cells. Only five bacterial isolates were obtained from industrial wastewater samples. Isolate NDSL2 withstood lead concentrations, MIC recorded at 180  $\mu\text{g/ml}$ , and selected as the most resistant isolate. By depending on morphological and molecular characterizations, it was identified as *Bacillus amyloliquefaciens* MPA 1034 strain. The optimized conditions for improving Pb-bioremoval efficiency were established. Maximum bioremoval of lead has been revealed 70% by growing and active bacterial cells after 3 days at pH 7, in the presence of lactose, sodium nitrate and yeast extract. SEM, EDX and FTIR analysis confirmed Pd uptake in and/or bound on bacterial cells. Thus, it could be recommended to use Pb-resistance bacterial strain as bioremoval and remediate polluted industrial effluents.

**Keywords:** bioremediation, bioremoval, *Bacillus*, wastewater, lead

### INTRODUCTION

Metals in general play important roles in the metabolism of microbial cells through acting as a cofactor for several enzymes. However, lead (Pb) is physiologically non-essential and causes health complications in animals and humans. Elevated lead concentrations can be toxic to microorganisms. Lead has several detrimental effects on microbial cells (Sun *et al.*, 2023). Some microbes possess lead detoxification mechanisms as reported by Mitra *et al.*, (2021).

Lead is one of the heavy metals which mainly derived from a variety of sources, one of these sources is the textile dyeing industry which has a negative significant impact on the environment. During the dyeing process, various metals are used as mordants e.g. lead white and lead chromate pigments. The dyeing process contributes up to one-fifth of water pollution and the textile dyeing industry is a major culprit. In particular in undeveloped countries, where weak regulations and enforcement prevail, wastewater from dye houses and garment factories is often directly dumped into rivers and streams. This discharge contains a toxic cocktail of carcinogenic chemicals, dyes, and heavy metals. Not only does it harm the environment, but it also contaminates essential drinking water sources (Khattab *et al.*, 2020 and Rujido-Santos *et al.*, 2022).

Nowadays, the use of microbes as remediators is a solution that is more environmentally compatible compared to chemical and physical approaches. In such polluted areas, microbes often use numerous methods to eliminate and detoxify heavy metals. Moreover, it is cheap, highly effective and not based on high technology. Until now, many microbial agents (bacteria, fungi, and microalgae) have been found as bioremediation approaches (Delangiz *et al.*, 2020).

Therefore, the present study aims to bioremediate lead through isolating lead-tolerant bacterial strain from industrial wastewater, improving bioremoval process, and investigating

the micromorphological and molecular changes to confirm lead adsorption and removal.

### MATERIALS AND METHODS

#### Source of wastewater

Samples of wastewater were obtained from various sites related to the textile industry. Five samples of effluent water were collected from Falcon, and El-Nabarawi factories at El-Mahala El-Kobra city, El-Garbia governorate, and should be mentioned here that these sampling sites are tanneries and spinning and weaving factories (Table 1). Concentrations of metal ions in all wastewater samples were analyzed in  $\mu\text{g/ml}$  according to Allen *et al.*, (1974). All samples were collected aseptically in sterile plastic containers and stored at 4°C for further use.

**Table 1. Source and description of wastewater samples**

No.	Sample description	Code
1.	The source of the sample was the first tank of the yarns dyeing process, El-Nabarawi factory.	NDF
2.	The source of the sample was the second tank of the yarns dyeing process, El-Nabarawi factory.	NDS
3.	The source of the sample was the first tank of the yarns dyeing process, Falcon factory.	FDL
4.	The source of the sample was the third tank of the yarns dyeing process, Falcon factory.	FDT
5.	The source of the sample was the industrial effluent resulting from the softening step, the Falcon factory.	FSS

#### Screening and isolation of Pb-resistance bacteria

The pour plate method was used for isolation. Under sterile conditions, one milliliter of wastewater samples was mixed with nutrient agar medium containing lead acetate, 100  $\mu\text{g/ml}$ . Petri plates were then incubated at 30°C for 48h. Colonies formed were picked up and purified.

#### Morphological and molecular identification of Pb-resistance bacterium

\* Corresponding author.

E-mail address: [eashour@mans.edu.eg](mailto:eashour@mans.edu.eg)

DOI: 10.21608/jacb.2024.281919.1079

The bacterial isolates were identified based on some cultural and morphological characteristics given in Bergey's Manual of Determinative Bacteriology (1994). In addition, the molecular identification of isolates was carried out by Sigma Scientific Services Company. Where, molecular and phylogenetic characterization of the isolates were performed based on 16S rRNA gene identification. The genomic DNA of the microbial isolates was extracted. The genomic DNA was used as a template for the amplification of 16S rRNA gene using universal primers. The PCR reaction consisted of 25 µL of PCR master mixture, 8 µL genomic DNA, 1 µL of the forward primer, 1 µL of the reverse primer, and 15 µL nuclease-free water. The obtained PCR sequences were used to find out the related sequences with known taxonomic information in the GenBank database (<http://www.ncbi.nlm.gov/blast>).

#### **Determination of minimum inhibitory concentration (MIC)**

The minimum inhibitory concentration (MIC) of lead for isolated bacteria was determined using the pour plate method (Aleem *et al.*, 2003; Ansari and Malik 2007). Nutrient agar plates supplemented with different concentrations (ranging from 100 to 350 µg/ml) of lead salt solutions were prepared. The inoculated plates were incubated at 30°C, followed by 72h until obvious growth of bacterial colonies was observed. The lowest concentration of the metal, which inhibits the growth was considered as the MIC of the metal against the tested isolate. Then, the most resistant isolate was selected for subsequent bioremoval experiments.

#### **Preparation of Inoculum**

Cell suspensions were prepared from fresh slants (48 h old). One milliliter from cell suspension was inoculated to 250 ml Erlenmeyer flask containing 50 ml sterile nutrient broth and incubated with shaking at 150 rpm at 30°C for 48h. Inoculums of bacterial cultures were adjusted at 10<sup>6</sup> cell/ml for further fermentation experiments.

#### **Lead bioremoval in shaking batch culture experiment**

Bioremoval of lead was assessed by growing the bacterial isolates interact with Pb(II) in Tryptone yeast extract broth medium at pH 7.2 (Ksheminska *et al.*, 2003). One mL of bacterial inoculum was inoculated in a 250 ml Erlenmeyer flask containing 50 ml of sterile broth medium mixed with lead, 100 µg/ml. Flasks were incubated at 30°C on shaking incubator. After the incubation period, the bacterial culture was centrifuged at 6000 rpm for 20 min. The supernatants were collected to determine the amount of the residual or unadsorbed Pb<sup>++</sup> using the atomic absorption spectrophotometer (AAS) according to Allen *et al.*, (1974). The bioremoval efficiencies were calculated using the following equation:

$$\% \text{ metals bioremoval} = [(C_i - C_r) \div C_i] \times 100$$

Where,

C<sub>i</sub> is the initial metal concentration in the growing medium (µg/ml), and C<sub>r</sub> is the residual metal concentration in the supernatant (µg/L).

#### **Optimization of Pb-bioremoval in shaking batch culture**

Some parameters were tested to reach the most suitable conditions of the bioremoval process for Pb (II). The performance of metal sorption by living microbial cells is influenced by environmental factors (incubation time and pH values) and essential nutritional factors (carbon and nitrogen sources). Data presented as bioremoval percentage.

#### **Time course**

Various incubation periods were tested from 1 to 4 days. One ml of bacterial inoculum (10<sup>6</sup> cell/ml) was cultured in a 250 ml Erlenmeyer flask containing 50 ml of sterile Tryptone yeast extract broth medium mixed with 100 µg metal salt /ml. All inoculated flasks were then incubated at

30°C in shaking incubator to determine the suitable time for metal bioremoval.

#### **Carbon source**

Glucose was replaced in tryptone yeast extract broth medium with different carbon sources; lactose, sucrose, maltose, fructose, and starch, at a concentration of 2%. Flasks were incubated for 2 and 3 days at 30°C with shaking at 150 rpm. The metal residual of each treatment was determined. Data presented as bioremoval percentage. The best carbon source was used in subsequent experiments.

#### **Nitrogen sources and their combinations**

Different nitrogen sources were examined in tryptone yeast extract broth medium contains the best carbon source and free of nitrogen. Nitrogen sources used were: NH<sub>4</sub>Cl<sub>2</sub>; (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>; (NH<sub>4</sub>)<sub>2</sub>PO<sub>4</sub>; NH<sub>4</sub>NO<sub>3</sub>; NO<sub>2</sub>NO<sub>3</sub>; yeast; beef extract and peptone at concentration 1%. Flasks were incubated for 2 and 3 days at 30°C with shaking at 150 rpm. The best nitrogen source was used in subsequent experiments. As for nitrogen sources combination, mixtures of the best organic and mineral nitrogen sources were: (1) 0.5 g NaNO<sub>3</sub>; (2) 0.5g beef; (3) 0.25g NaNO<sub>3</sub>+0.25g beef; (4) 0.125g NaNO<sub>3</sub>+0.375g beef; (5) 0.375g NaNO<sub>3</sub>+0.125g beef. Flasks were incubated for 2 and 3 days at 30°C with shaking at 150 rpm. The metal residual of each treatment was determined. Data presented as bioremoval percentage.

#### **Hydrogen ions concentration (pH values)**

Various pH values from 5 to 9 were tested for their effect on the bioremoval activity of the tested bacterial cells. One ml of bacterial inoculum (10<sup>6</sup> cell/ml) was cultured in a 250 ml Erlenmeyer flask containing 50 ml of sterile Tryptone yeast extract broth medium mixed with lead metal (100 µg /ml). Then, flasks were incubated for 2 and 3 days at 30°C with shaking at 150 rpm.

#### **Characterizations**

The bacterial cells grown in optimized tryptone yeast extract broth medium supplemented with and without (control) lead solution (100 µg/ml) were centrifuged and cell pellets were used for SEM, EDX and FTIR analysis.

#### **Scanning electron microscopy (SEM)**

SEM technique reveals detailed morphological characterization and changes in bacterial cells. The prepared pellets were observed in a Jeol JSM-6510 LV SEM, operating at 30 kV at the Electron Microscopy Unit, Faculty of Agriculture, Mansoura University, Egypt.

#### **Energy dispersive X-ray spectroscopy (EDX)**

EDX techniques allow for detailed elemental analysis of bacterial samples. A SEM equipped with EDX was used for analyzing the specimens (Cairns Advanced Analytical Centre, 2017). EDX spectra were used to identify the presence of specific elements within the microbial cells and their distribution.

#### **Fourier Transform infrared spectroscopy (FTIR)**

FTIR was used to investigate the various functional groups involved in bioremoval of lead and possible metal binding sites of functional groups present in the biomass of the tested isolate. FTIR spectroscopy provides valuable information about the biochemical composition and structural properties of bacterial cells. The prepared sample was placed into the FTIR spectrometer "ThermoFisher Scientific" at the Spectral Analysis Unit, Faculty of Science, Mansoura University, Egypt. Infrared radiation passed through the sample, and the resulting infrared spectrum (within the range of 500-4000 cm<sup>-1</sup> spectrum) was recorded. Spectral data were analyzed using software to identify characteristic peaks and

patterns associated with microbial biomolecules such as proteins, lipids, carbohydrates, and nucleic acids.

**Heavy metals analysis**

The residual heavy metal ions concentration of the microbial samples was detected in the supernatants and analyzed using a Buck Scientific Accusys 211 series “Atomic Absorption Spectrophotometer” (58 Fort Point St. East Norwalk, CT 06855, USA) in the Atomic Absorption Lab, Unit of Genetic Engineering and Biotechnology, at the Faculty of Sciences, Mansoura University, by an air/acetylene flame system. The concentration of lead in the samples was determined in µg/ml (Allen *et al.*, 1974).

**RESULTS AND DISCUSSION**

Data represented in Table (2) indicate metal content in wastewater samples thrown from textile and dyeing factories. Among the detected metals, the wastewater samples contain lead in various amounts.

**Table 2. Initial concentrations of detected heavy metals (µg/ml) of five wastewater samples from tanneries and spinning and weaving factories.**

Metals detected	NDF	NDS	FDf	FDT	ESS
Co <sup>++</sup>	0.067	0.023	0.059	0.029	0.019
Pb <sup>++</sup>	0.140	0.009	0.102	0.008	0.025
Ni <sup>++</sup>	0.728	0.019	0.498	0.019	0.012
Zn <sup>++</sup>	0.012	0.028	0.048	0.024	0.027
Cd <sup>++</sup>	0.041	0.014	0.034	0.013	0.023

**Isolation and screening of lead-resistant bacteria**

From wastewater samples, only five bacterial isolates were obtained that were able to grow on nutrient agar medium supplemented with lead acetate (100 µg/ml). Then, the effect of several concentrations of Pb<sup>++</sup> from 100 to 200 µg/ml was tested to determine MICs. Data represented in Table (3) show the MICs of metal (Pb<sup>++</sup>) on the bacterial growth where their response to lead toxicity was varied. Isolates NDFL1 and NDSL2 were high Pb- resistance bacteria. Isolate NDSL2 withstood lead concentrations up to 180 µg/ml, which is considered the most Pb-resistant one.

**Table 3. Minimum inhibitory concentration (MIC) of lead on isolated bacteria**

Metal concn. (µg/ml)	Bacterial isolates				
	NDFL1	NDSL2	FDTL3	FDL1	FSSL5
100	+	+	+	+	+
110	+	+	+	+	+
120	+	+	+	+	+
130	+	+	+	+	-
140	+	+	+	+	-
150	+	+	-	+	-
160	+	+	-	-	-
170	+	+	-	-	-
180	-	+	-	-	-
190	-	-	-	-	-
200	-	-	-	-	-
Mic (µg/ml)	170	180	140	150	120
(+) Growth	(-) No growth				

Many previous studies confirm bacterial tolerance against lead, Al-Ansari *et al* (2021) showed high results compared to the present values of MIC where the MIC of lead for *P. aeruginosa* strain RA-14 was 2800 µg/ml, and Babar *et al.*,(2021) also reported that the MIC of Pb<sup>++</sup> for *Bacillus altitudinis* MT422188 was 5 mM. In addition, Mitra *et al.*,(2021) reported that microbes inhabiting Pb-contaminated areas are found to have evolved distinctive mechanisms to successfully thrive in the Pb-contaminated environment


without exhibiting any negative effects on their growth and metabolism.

Depending on the obtained results, the highest Pb-resistant bacterial isolate was selected for the optimization of metal bioremoval experiments.

**Characterization and identification of bacterial isolate**


Identification of the most resistant isolate, NDSL2, was depended on morphological and molecular characterizations.

Morphological characterization: On nutrient agar medium after 48 h, a culture of NDSL2 showed a gray smooth colony with an irregular shape. Rod cells are arranged mostly in single cells and sometimes in diploe, cells length between 2-2.4 µm, gram-positive and spore formers were obviously in the microscopic examination (Fig. 1).



**Fig. 1. Microscopic photograph of Pb-resistant isolate, NDSL2, showing morphological characterization**

**Molecular characterization:** The molecular identification of Pb-resistant isolate, NDSL2, was carried out by Sigma Scientific Services Co. The obtained sequencing of the 16S rRNA gene of the bacterial isolate, NDSL2, was compared with the GenBank’ deposited sequences and revealed that the isolate is almost *Bacillus* sp. and showed 99.88% similarity with *Bacillus amyloliquefaciens* MPA 1034 strain. The phylogenetic relationship of the identified *Bacillus* sp. and other similar strains can be shown in Fig. (2).



**Fig. 2. Phylogenetic tree constructed using the 16S rRNA sequence of Bacillus amyloliquefaciens MPA 1034 strain and related strains using NCBI BLAST analysis and MEGA6 software.**

**Optimization of metal bioremoval in shacking batch culture**

Metal removal by living cells is growth-dependent and may be altered by nutritional and environmental

conditions that influence microbe production. Thus, to improve lead removal by the resistant microbial strain, *B. amyloliquefaciens* MPA 1034, some environmental (contact time and pH) and nutritional (carbon and nitrogen) factors were tested.

### Time course of metal bioremoval

The incubation period has a significant impact on the bioremoval process as it is directly affecting the microbial growth and their subsequent contact and uptake of the metals. Data illustrated in Fig. (3) indicate the bioremoval of Pb<sup>++</sup> by *B. amyloliquefaciens* MPA 1034 through different incubation periods from 1 to 4 days. It was found the maximum lead removal reached 34.9 and 37.7 % after 2 and 3 days, respectively. These results are in harmony with those obtained by Salman (2012) who reported the effect of incubation periods on Pb<sup>++</sup> biosorption by *Brevundimona svesicularis* (C18), the highest uptake was recorded after 60h, reaching 30.75% of the initial metal concentration. Also the maximum peak of uptake capacity of Pb<sup>++</sup> was obtained after 48h of incubation period reaching 34.68% by *Bacillus circulans* (D21) and 49.82% after 36h by *Klebsiella mobilis* (C19).




Fig. 3. Effect of incubation time on lead removal by *B. amyloliquefaciens* MPA 1034

### Effect of carbon source

Data in Figs. (4) show the effect of carbon sources on the bioremoval of Pb<sup>++</sup> by *Bacillus amyloliquefaciens* MPA. Replacing glucose with fructose and maltose caused high decreases in lead bioremoval by *B. amyloliquefaciens* MPA, while enrichment of medium with sucrose and lactose resulted in a high bioremoval. Lactose was the best carbon source for lead bioremoval, which achieved 48.9 and 49.3% after 2 and 3 days respectively.




Fig. 3. Effect of carbon sources on lead removal by *B. amyloliquefaciens* MPA

Carbon is considered the main nutrient element in bacteria metabolism, which bacteria use it build up their biomass as anabolic reactions (Rittmann and Mc Carty,

2001). The properties of the organic carbon source play an important role in the operation of biological processes. The assessment has been made mostly by the conventional analyses for proteins and polysaccharides (Li and Yang, 2007; Ye et al.,2011; Zhao et al.,2018), which affect the bacterial growth, and reflect on the ability of the microorganisms for heavy metal bioremoval.

### Effect of nitrogen sources

Lead removal by *B. amyloliquefaciens* MPA was affected positively and negatively with nitrogen sources, whereas ammonium sulphate was the best mineral nitrogen source which recorded 50%, while yeast extract was the best organic nitrogen source which recorded 51.3 % after 2 days of incubation (Fig. 5).




Fig. 5. Effect of nitrogen sources on lead by *Bacillus amyloliquefaciens* MPA .

### Effect of the mix of best mineral and organic nitrogen sources

The best mineral and organic nitrogen sources were tested in different proportions (by quarters) to determine the best mixture for *B. amyloliquefaciens* MPA to enhance the bioremoval of the heavy metals. Using mixtures from sodium nitrate and yeast extract caused an increase in the bioremoval of lead by *B. amyloliquefaciens* MPA, which achieved the highest lead bioremoval 67.4 % after 2 days by using a mixture of 50% sodium nitrate (mineral nitrogen source) and 50% of yeast extract as the organic nitrogen source in Fig. (6).





Fig. 6. Effect of mineral and organic nitrogen sources ratios on the bioremoval of lead by *B. amyloliquefaciens* MPA

### Effect of pH

One of the important factors to consider in the growth of many microorganisms is the pH, since it plays an important role in how microorganisms will interact with the environment (Mougi 2023). Herein, data illustrated in Fig. (7) revealed the wide range of pH in which *Bacillus amyloliquefaciens* MPA could grow and has a high potential for lead bioremoval. Lead bioremoval by *Bacillus amyloliquefaciens* MPA was high efficient in the pH range

from 6.5 to 8.0. Although the highest lead bioremoval was achieved at pH 7 and 7.5, which recorded lead bioremoval 70.3% at pH 7 as well as 70.5 at pH 7.5 after 3 days. Raising the initial pH over 8.0 decreased lead bioremoval, which also happened when lowering pH below 6.



**Fig. 7. Effect of initial medium pH on the bioremoval of lead by *B. amyloliquefaciens* MPA**

The degree of bioremoval efficiency of microbial cells for heavy metal ions depends upon various external operating factors such as pH, and contact time, in addition to the aqueous nutritional environment (Aryal and Liakopoulou-Kyriakides 2015).

The pH is a major factor that affects the solution chemistry of metal ions and the surface functional groups of the microbial cell wall, where H<sup>+</sup> ions replace some of the positive ions from the biomass surface. Bioremoval of the capacity of metal cations also increases with rising in values of pH, that is, may be due to the more negative binding sites exposed on biomass surface. On the contrary, at low pH values, the binding sites of the cell wall are blocked and associated with hydrogen ions that prevent the access of metal cations (Singh et al. 2022).


**Scanning electron microscopy (SEM) and Energy Dispersive X-ray (EDX)**

SEM gives a visualization of the microbial cell's morphology at high magnifications, and these images can be used to identify and define changes in the shape, size and

surface of the microbial cells after heavy metal exposure. In addition, the EDX spectrum perform the elemental mapping of metal loaded by the surface of microbial cells.

Comparing SEM images of untreated *Bacillus amyloliquefaciens* MPA biomass cultured for 48 in optimized tryptone yeast extract broth medium and Pb-treated (100 µg/ml) and SEM images shown in Fig. (8 a&b). It was observed that untreated *B. amyloliquefaciens* MPA cells showed regular long rods (Fig. 8a). While cell morphology of Pd(II) treated revealed many cells in dissimilar size and deformation of the cells due to wrinkle of cell wall (Fig. 8b). These images were further confirmed by EDX analysis for the elemental mapping of metal loaded on *B. amyloliquefaciens* MPA ' cells and it was observed metal ions distribution on the surface of the bacterial cells (Fig. 8 a&b). Along with the corresponding metal ions, it was found that C and O are major elements present in both untreated and Pb-treated cells of *B. amyloliquefaciens* MPA. The EDX spectrum of lead-loaded *B. amyloliquefaciens* MPA appeared the peak of lead, 0.06%. Consequently, the existence of Pb(II) ion on the cell surface of *B. amyloliquefaciens* MPA was confirmed by the EDX spectrum.

Analysis with SEM and EDX in previous studies confirmed the adsorption of metals on the bacterial biomass. For instance, SEM images of Gram-positive bacteria *Bacillus cereus* FIT10 (Dhanwal et al.,2018) and *Staphylococcus hominis* strain AMB-2 (Rahman et al.,2019) showed clear adsorption of Pb after heavy metals treatment. Moreover, EDX analysis also confirmed the presence of a Pb signal on the bacterial surface. In addition, Shao et al.,(2019) investigated extracellular accumulation by SEM with EDX and found that extracellular accumulated Pb reached 61.7–95.9% of the total accumulation. However, when they examined the morphological differences of *Bacillus* sp. MPR-3 cells in the presence and absence of lead ions by SEM. They demonstrated that bacterial cells became bloated, and smooth, and formed precipitates in the presence of Pb. The distinct morphological alterations, deformation, and severe membrane damage prove Pb toxicity to bacteria that result in different degrees of damage.



**Fig. 8. Scanning electron micrograph (arrow points to changes) and EDX spectrum (arrow points to Pb peak) of *Bacillus amyloliquefaciens* MPA biomass cultured in optimized tryptone yeast extract broth medium. A) control – absence of metal; B) treated with lead solution (100 µg/ml).**


**Fourier Transform Infrared Spectroscopy (FTIR)**

The FTIR analyses were run to investigate the characteristic functional groups on the surface of the microbial

cell and identify the change that arises from the complex formation after metal exposure. The characteristic functional groups may have crossed a shift in their values than the control

samples or disappearance of the frequency of a definite group. Here, the scale of the verified frequencies was expressed as the wavenumber in the ranges of 500 to 4000  $\text{cm}^{-1}$ .

Cells of *Bacillus amyloliquefaciens* MPA that were cultured in optimized tryptone yeast extract broth medium (both lead-free and in the presence of lead solution, 100  $\mu\text{g}/\text{ml}$ ) were characterized by infrared spectroscopic analyses. The results stated the representative frequencies and the deduced functional groups are shown in Fig. (9). The absorption band at  $\nu=1742\text{ cm}^{-1}$  for strong stretching C=O “esters” of the control sample was absent in the analysis of the cells treated with lead. On the contrary, a new absorption band at 1232 is attributed to strong stretching C-O or medium stretching C-N “amine” groups that appeared in the FTIR spectrum of cells treated with lead. Furthermore, the stretching absorption band is related to the formation of a new bond between lead and oxygen atom “Pb-O” at  $\nu=618\text{ cm}^{-1}$ .



**Fig. 9. FTIR spectra of *Bacillus amyloliquefaciens* MPA ' biomass cultured in optimized tryptone yeast extract broth medium A) Lead-free (control) and B) in the presence of lead solution, (100  $\mu\text{g}/\text{ml}$ ).**

FTIR spectroscopy is a powerful tool for monitoring and revealing microorganism composition and responses to the environment. So, it was performed to confirm the interactions between metal ions and functional groups on the surface of the microbial cells in comparison to untreated ones (Dhanwal et al., 2018; Rizvi and Saghir-Khan 2019; Pagnucco et al., 2023 and Tiquia-Arashiro 2023). Where, Dhanwal et al., (2018) identified the surface chemical functional groups of the metal tolerant strain, *Bacillus cereus* FIT10, by FTIR spectroscopy as hydroxyl, carboxyl, amine, and halide, which might be involved in the biosorption of heavy metals, lead, chromium, copper, and nickel.

## REFERENCES

- Al-Ansari M.M., Benabdelkamel H., AlMalki R.H., Abdel Rahman A.M. Alnahmi E., Masood A., Ilavenil S. and Choi K.C. (2021). Effective removal of heavy metals from industrial effluent wastewater by a multi metal and drug resistant *Pseudomonas aeruginosa* strain RA-14 using integrated sequencing batch reactor. *Environ. Res.*, 199:111240.
- Aleem A., Isar J. and Malik (2003). Impact of long-term application of industrial wastewater on the emergence of resistance traits in *Azotobacter chroococcum* isolated from rhizospheric soil. *Bioresour. Bioresource Technol.*, 86(1):7-13.
- Allen S., Grimshaw H.M., Parkinson J.A. and Quarmby C. (1974). *Chemical Analysis of Ecological Material*, Oxford, Blackwell Scientific Publication., p. 521.
- Ansari M.I. and Malik A. (2007). Biosorption of nickel and cadmium by metal resistant bacterial isolates from agricultural soil irrigated with industrial wastewater. *Bioresour. Technol.*, 98: 3149-3153.
- Aryal M. and Liakopoulou-Kyriakides M. (2015). Bioremoval of heavy metals by bacterial biomass. *Environ. Monit. Assess.*, 187(1):4173. DOI: 10.1007/s10661-014-4173-z.
- Babar Z., Khan M., Chotana G.A., Murtaza G. and Shamim S. (2021). Evaluation of the potential role of *Bacillus altitudinis* MT422188 in nickel bioremediation from contaminated industrial effluents. *Sustainability.*, 13:7353. <https://doi.org/10.3390/su13137353>.
- Bergey D. (1994). *Bergey's Manuals of Determinative Bacteriology*. In: Holt John G., eds. Lippincott Williams and Wilkins.
- Cairns Advanced Analytical Centre (2017). *Biological Sample Preparation for the SEM*. <https://research.jcu.edu.au/archive/enabling/aac/cairns-aac/biological-sample-preparation-for-the-sem>
- Delangiz N., Varjovi M.B., Lajayer B.A. and Ghorbanpour M. (2020). Beneficial microorganisms in the remediation of heavy metals. *Molecular aspects of plant beneficial microbes in agriculture*. Elsevier Inc. <https://doi.org/10.1016/B978-0-12-818469-1.00034-1>.
- Dhanwal P., Shruti A., Dudeja S., Badgajar H., Chauhan R., Kumar A., Dhull P., Chhokar V., Beniwal V. (2018). Biosorption of heavy metals from aqueous solution by bacteria isolated from contaminated soil. *Water Environ. Res.*, 90:424-430. DOI.org/10.2175/106143017X15131012152979
- Khatab T.A., Abdelrahma M.S. and Rehan M. (2020). Textile dyeing industry: environmental impacts and remediation. *Environ. Sci. Pollut. Res.*, 27:3803–3818. <https://doi.org/10.1007/s11356-019-07137-z>
- Ksheminska H., Jaglarz A., Fedorovych D., Babyak L., Yanovych D., Kaszycki P. and Koloczek H. (2003). Bioremediation of chromium by the yeast *Pichia guilliermondii*: toxicity and accumulation of Cr(III) and Cr(VI) and the influence of riboflavin on Cr(III) tolerance. *Microbiol. Res.*, 158:59–67. <https://doi.org/10.1078/0944-5013-00178>.

- Li X.Y. and Yang S.F. (2007). Influence of loosely bound extracellular polymeric substances (EPS) on the flocculation, sedimentation and dewater ability of activated sludge. *Water Res.*, 41:1022–1030.
- Mitra A., Chatterjee S., Katakai S., Rastogi R., and Gupta D. (2021). Bacterial tolerance strategies against lead toxicity and their relevance in bioremediation application. *Environ. Sci. and Pollution Res.*, 28:10.1007/s11356-021-12583-9.
- Mougi A. (2023). Eco-evolutionary dynamics in microbial interactions. *Scientific Reports*, 3:13(1):9042. DOI: 10.1038/s41598-023-36221-1.
- Pagnucco G., Overfield D., Chamlee Y., Shuler C., Kassem A., Opara S., Najaf H., Abbas L., Coutinho O., Fortuna A., Sulaiman F., Farinas J., Schittenhelm R., Catalfano B., Li X. and Tiquia-Arashi S.M. (2023). Metal tolerance and biosorption capacities of bacterial strains isolated from an urban watershed. *Front. Microbiol.* 14:1278886. DOI:10.3389/fmicb.2023.1278886
- Rahman Z., Thomas L. and Ved Pal Singh V.P. (2019). Biosorption of heavy metals by a lead (Pb) resistant bacterium, *Staphylococcus hominis* strain AMB-2. *J. Basic Microbiol.*, 59:477-486. DOI: 10.1002/jobm.201900024
- Rittmann B.E. and McCarty P.L. (2001). *Environmental Biotechnology: Principles and Applications*. McGraw-Hill.
- Rizvi A. and Saghir-Khan M. (2019). Putative role of bacterial biosorbent in metal sequestration revealed by SEM-EDX and FTIR. *Indian J Microbiol.*, 59(2):246-249. Doi: 10.1007/s12088-019-00780-7.
- Rujido-Santos I., Herbelo-Hermelo P., Barciela-Alonso M.C., Bermejo-Barrera P. and Moreda-Piñeiro A. (2022). Metal Content in Textile and (Nano)Textile Products. *Internat. J. of Environ. Res. and Public Health*, 19(2):944. <https://doi.org/10.3390/ijerph19020944>
- Salman B.E.A. (2012). Use of microorganisms for bioremoval of heavy metals from some polluted sources. Ph.D. Faculty of Agriculture. Zagazig University.
- Shao H., Qi Y., Liang S., Qin S. and Yu J. (2019). Polypropylene composite hollow fiber ultrafiltration membranes with an acrylic hydrogel surface by in situ ultrasonic wave-assisted polymerization for dye removal. *J. Appl. Polym. Sci.*, 136:1-10. <https://doi.org/10.1002/app.47099>.
- Singh N., Goutam U. and Ghosh M. (2022). Deep-marine bacteria-the Frontier alternative for heavy metals bioremediation. *Development in wastewater treatment research and processes*. Elsevier, Amsterdam, pp 429-450.
- Sun X., Sun M. and Chao Y. *et al.* (2023). Effects of lead pollution on soil microbial community diversity and biomass and on invertase activity. *Soil Ecol. Lett.*, 5:118–127. <https://doi.org/10.1007/s42832-022-0134-6>
- Tiquia-Arashi S, Li X, Pokhrel K, Kassem A, Abbas L, Coutinho O, Kasperek D, Najaf H and Opara S (2023). Applications of Fourier Transform-Infrared spectroscopy in microbial cell biology and environmental microbiology: advances, challenges, and future perspectives. *Front. Microbiol.*, 14:1304081. Doi:10.3389/fmicb.2023.1304081
- Ye F., Peng G. and Li Y. (2011). Influences of influent carbon source on extracellular polymeric substances (EPS) and physicochemical properties of activated sludge. *Chemosphere.*, 84: 1250–1255.
- Zhao J., Li Y., Chen X. and Li Y. (2018). Effects of carbon sources on sludge performance and microbial community for 4-chlorophenol wastewater treatment in sequencing batch reactors. *Bioresour. Technol.*, 255:22–28.

## الإزالة البيولوجية للرصاص بواسطة *Bacillus amyloliquefaciens* MPA 1034 المعزولة من مياه الصرف الصحي في مصانع النسيج بالمحلة الكبرى، مصر

مي محمود احمد الجوجري ، محمد عبدالله العوضى سليم و إيمان حسين عاشور

قسم الميكروبيولوجي، كلية الزراعة، جامعة المنصورة

### المخلص

تعتبر المعادن الثقيلة أخطر الملوثات البيئية حيث أدى النشاط الصناعي إلى إطلاق كميات كبيرة من النفايات السائلة الضارة والتي تحتوي على المعادن الثقيلة السامة في البيئة، ولذلك هناك حاجة ماسة وملحة لمعالجة مياه الصرف الصناعي. تهدف هذه الدراسة إلى الإزالة الحيوية لعنصر الرصاص بواسطة الخلايا البكتيرية النشطة والمقاومة للرصاص. تم الحصول على خمس عزلات بكتيرية فقط من عينات مياه الصرف الصناعي، وكانت العزلة NDSL2 مقاومة لتركيزات الرصاص وسجلت MIC عند 180 ميكروجرام/مل وتم اختيارها كأكثر عزلة مقاومة. وعلى أساس الخصائص المورفولوجية والجزيئية، تم تحديدها على أنها سلالة *Bacillus amyloliquefaciens* MPA 1034. وقد تم الوصول إلى الظروف المثلى لتحسين كفاءة الإزالة الحيوية للرصاص. وسجل الحد الأقصى 70% من الإزالة الحيوية للرصاص بواسطة الخلايا البكتيرية النشطة بعد 3 أيام وعند درجة حموضة 7، وفي وجود اللاكتوز و نترات الصوديوم ومستخلص الخميرة. أكد تحليل SEM و EDX و FTIR إدمصاص أو امتصاص (أو كلاهما) معدن الرصاص بالخلايا البكتيرية. وبالتالي يمكن التوصية باستخدام هذه السلالة البكتيرية في الإزالة الحيوية للرصاص ومعالجة النفايات السائلة الصناعية الملوثة.